

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com

2. PASTURES, SEED CROPS & TURF (continued)

Situation & Crop	Weeds	State	Rate	Critical Comments
Pastures - High Volume Spraying Technique	Caltrop, Capeweed, Chickweed, Mustard, Paterson's Curse, Shepherd's Purse, Saffron, Slender, Spill or Variegated Thistle, Turf Weed, Wild Radish, Wild Turnip	NSW, ACT, Vic only	350 mL/1.4 Lha	Spray actively growing 6-8 weeks old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking with at least 5 times heavier than normal until weeds have been reduced but before arrival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. Refer 'Legume Tolerance'.
	Refer to Weed Table	WA only	825 mL/1.85 Lha	Refer 'Legume Tolerance'.
Capeweed, Dandelion, Seeding Dock, Three Corners, Jack, Geranium, Erodium, Mustard, Annual Thistles, Turf Weed, Wild Radish, Wild Turnip	Refer to Weed Table	WA only	500 mL/ha	PRECAUTION: An increased quantity of potassium salts may be eaten by stock using Spray/Graze, eg. Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's Curse. Observe Withholding Period of at least 7 days after spraying.
	Refer to Weed Table	SA only	700 mL/ha	Refer 'Legume Tolerance'.
Turf	Refer to Weed Table	All States	750 mL/ha or 5 L/ha per 100 sq metres	Newly laid - Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See 'Legume Tolerance'. Buffalo grass is more easily damaged than other grasses.
	Refer to Weed Table	All States	1425 L/ha or 1425 mL per 100 sq metres	Established - Apply in high volume to actively growing weeds. Do not mow for 2 days before application. Some transitory damage may occur to the turf grasses and strawberry clover. See 'Legume Tolerance'. Buffalo grass is more easily damaged than other grasses.

3. SPOT SPRAYING

Situation & Crop	Weeds	State	Mixing Rates/Critical Comments
High Volume Spraying	Refer to Weed Table	All States	Add 1/10th of rate on Weed Table to 150 L of water. Each 100 L of mix will cover 1000 m ² (1/100th ha) eg. If rate in Weed Table is 1 L/L use 140 mL/150 L water.
Knapsack Application	Refer to Weed Table	All States	Add 1/100th of rate on Weed Table to 10 L of water. Each 10 L of mix will cover 100 m ² (1/100th ha) eg. If rate in Weed Table is 1 L/L use 14 mL/10 L water.

4. Non Crop Areas, Fallow Land, Industrial & Commercial Areas, Rights-of-Way

Situation & Crop	Weeds	State	Rate	Critical Comments
Non Crop Areas, Fallow Land, Industrial & Commercial Areas, Rights of Way	Refer to Weed Table	All States	Refer to Weed Table	Apply at rates shown in Weed Table when weeds only are present and no crop or pasture is present.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: CEREALS, PASTURE: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application.

APPLICATION/MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

BOOM SPRAYING - Use 30-120 L water per hectare. - Use minimum of 170 L water per hectare for LIKESD.

AERIAL SPRAYING - Use 10-90 L water per hectare.

RESISTANT WEEDS WARNING **GROUP 1 HERBICIDE**

Sanonda MCPA 500 Selective Herbicide (MCPA 500) is a member of the Phenoxys group of herbicides. MCPA 500 has the Disruptors of plant cell growth mode of action. For weed resistance management MCPA 500 is a Group 1 herbicide. Some naturally-occurring weed biotypes resistant to MCPA 500 and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by MCPA 500 or other disruptors of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Sanonda Australia Pty Ltd accepts no liability for any losses that may result from the failure of MCPA 500 to control resistant weeds.

EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with commercially available tank and equipment cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

COMPATIBILITY/TANK MIXES

This product can be tank mixed with Agility, Aity/Associate, Kambari, Bromicide, Flowable Diuron, Archer, Serfoni, Spraysave, Flowable Terbutryn, Tordon and Triburon. The effectiveness of Comrad, and Dimetholac may be reduced by alkaline hydrolysis if tank mixed with this product. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

1. FIELD CROPS (continued)

Situation & Crop	Weeds	State	Rate	Critical Comments	
Field peas	Procky Lettuce (Lactuca serriola)	NSW, Vic, Tas, SA only	4 leaf	125 mL plus 125 mL Nufarm Agly Selective Herbicide	FIELD PEAS: Apply early post-emergence after the first node stage and before the start of flowering. FIELD PEAS: Apply early post-emergence after the first node stage and before the start of flowering. Field peas grown on high pH soils in the presence of free lime may be less tolerant to Agly Agly selective herbicide. APPLICATION AND WEED CONTROL: Apply when weeds are actively growing. For optimum results apply 4 to 6 weeks post-sowing. Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: - high crop and weed density - abnormal weed growth (including early flowering), high rate of product up to the maximum rate of application specified for that weed may be required. Nufarm Agly Selective Herbicide will not effectively control: - regrowth of suppressed weeds, - regrowth from rhizomes or roots, - weeds growing under stress from previous herbicide applications.
		NSW, ACT only	4 leaf	125 mL plus 125 mL Nufarm Agly Selective Herbicide	
		NSW, ACT, Vic, SA, Tas only	6 leaf	150 mL plus 150 mL Nufarm Agly Selective Herbicide	
		NSW, ACT, Vic, SA, Tas only	6 leaf	150 mL plus 150 mL Nufarm Agly Selective Herbicide	
Charlock (Wild mustard) (Sinapis arvensis)	Refer to Weed Table	Old only	300 mL/2.1 Lha	Apply when crop is 10-15 cm tall with at least 170 L/ha of water. Do not spray after bud set. Some white effect may be seen following application but crop usually recovers quickly. Do not spray by aircraft.	
		NSW, ACT, Vic, SA, Tas only	300 mL/1.1 Lha		

Situation & Crop

Situation & Crop	Weeds	State	Rate	Critical Comments
Lusweed	Refer to Weed Table	Old only	300 mL/2.1 Lha	Apply when crop is 10-15 cm tall with at least 170 L/ha of water. Do not spray after bud set. Some white effect may be seen following application but crop usually recovers quickly. Do not spray by aircraft.
		NSW, ACT, Vic, SA, Tas only	300 mL/1.1 Lha	

Sugar Cane

Situation & Crop	Weeds	State	Rate	Critical Comments
Blue Top, Chinese Burr, Fanleaf Weed, Gamba Pea, Bell Vine, Streaked Rattle Pod, Bromesec, Pook Convolvulus, Cupids Flower, Merremia Vine, Morning Glory	Refer to Weed Table	Old only	1.4 L/ha	POST-EMERGENT: Apply as directed inter-row spray. DO NOT exceed rates with phenox sensitive varieties.
		NSW, ACT, Vic, WA only	3 L/ha	Apply in Autumn when Hoarhound plants are at the seedling stage. If crop cover or medic plants are present they will be damaged by the application.

2. PASTURES, SEED CROPS & TURF (continued)

Situation & Crop	Weeds	State	Rate	Critical Comments
Grass Pasture	Refer to Weed Table	All States	700 mL/4 Lha	Apply to established grass based pastures. Damage may occur to Legumes if present. See 'Legume Tolerance' below. NOT to be used on Pasture's Curse and Variegated Thistle may be poisonous to stock after spraying.
		SA only	1.1 L/ha 400 mL/ha	Use this mixture only if spraying is delayed until early winter. Rosettes should not exceed 20 cm diameter.
Sodier Thistle	Refer to Weed Table	All States	700 mL/2.1 Lha	Use low rate on seedlings with no more than 2 true leaves. Use 100 L of water for best results. If clover present do not exceed 600 mL.
		NSW, ACT, Vic, WA only	3 L/ha	Apply in Autumn when Hoarhound plants are at the seedling stage. If crop cover or medic plants are present they will be damaged by the application.
Grass Seed Crops	Refer to Weed Table	Old, SA, Tas, Vic, WA only	700 mL/4 Lha	Apply to established grass seed crops.
		NSW, ACT, Vic, WA only	700 mL/1.5 Lha	Do not exceed rate specified. Spray in Autumn and early Winter.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LIFT HERE

Sanonda
MCPA 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L MCPA
present as the dimethylamine salt

GROUP 1 HERBICIDE

For the selective control of broadleaf weeds in cereals, inseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE USING PRODUCT

APVMA Approval No.: 63433/1209

SANONDA (Australia) Pty Ltd
Suite 822, 51 Koola Road Towns,
Queens Road, Melbourne, VIC 3004
Website: www.sanonda.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.com